

EXTENSOR POLLICIS LONGUS TENDINITIS



■ ■ ■ Description

Extensor pollicis longus tendinitis is characterized by pain in the back of the wrist due to inflammation of one of the tendon sheaths (lining) or strain of one of the tendons to the thumb, the extensor pollicis longus (EPL). The lining secretes a fluid that lubricates the tendon. When the lining becomes inflamed, the tendon cannot glide smoothly in its covering (sheath). The EPL tendon is the anchor of the EPL muscle, which is important for straightening the thumb. This may be a grade 1 or 2 strain of the tendon. A *grade 1 strain* is a mild strain. There is a slight pull without obvious tearing (it is microscopic tendon tearing). There is no loss of strength, and the tendon is the correct length. A *grade 2 strain* is a moderate strain. There is tearing of tendon fibers within the substance of the tendon or where the tendon meets the bone or muscle. The length of the whole muscle-tendon-bone unit is increased, and there is usually decreased strength. A *grade 3 strain* is a complete rupture of the tendon.

■ ■ ■ Common Signs and Symptoms

- Vague, diffuse pain and tenderness over the back of the wrist
- Pain that is worse with straightening the thumb
- Locking, catching, or triggering of the thumb joint
- Limited motion of the thumb
- Little or no swelling, warmth, or redness of the back of the wrist
- Crepitation (a crackling sound) when the tendon or wrist is moved or touched

■ ■ ■ Causes

- EPL tendinitis becomes more likely with repeated injury to the thumb muscle-tendon units and with repetitive motions of the hand and wrist due to friction of the tendon within the sheath (lining)
- Sudden increase in activity or change in activity

■ ■ ■ Risk Increases With

- Sports that involve repetitive hand and wrist motions, including tennis, golf, and bowling
- Previous wrist fracture
- Heavy labor
- Poor physical conditioning (strength and flexibility)
- Inadequate warm-up before practice or play
- Previous wrist immobilization or other injury

■ ■ ■ Preventive Measures

- Appropriately warm up and stretch before practice or competition.
- Allow time for adequate rest and recovery between practices and competition.

- Maintain appropriate conditioning:
 - Forearm, wrist, and hand flexibility
 - Muscle strength and endurance
- Use proper technique.

■ ■ ■ Expected Outcome

This condition is usually curable within 6 weeks if treated appropriately with conservative treatment and resting of the affected area.

■ ■ ■ Possible Complications

- Prolonged healing time if not appropriately treated or if not given adequate time to heal
- Chronically inflamed tendon, causing persistent pain with activity that may progress to constant pain, restriction of motion of the tendon within the sheath, and potentially rupture of the tendon
- Recurrence of symptoms, especially if activity is resumed too soon
- Risks of surgery, including infection, bleeding, injury to nerves (numbness of the thumb), continued pain, incomplete release of the tendon sheath, recurrence of symptoms, cutting of the tendon, thumb weakness, thumb stiffness, and inability to straighten the thumb

■ ■ ■ General Treatment Considerations

Initial treatment consists of medication and ice to relieve the pain and reduce the inflammation, stretching and strengthening exercises, and modification of the activity that initially caused the problem. These all can be carried out at home, although referral to a physical therapist or athletic trainer may be recommended. Occasionally a cast, brace, or splint may be prescribed to reduce motion, helping alleviate inflammation. An injection of cortisone to the area around the tendon is often attempted. Surgery to release the inflamed tendon lining may be needed. If the tendon is ruptured, the tendon will require surgical repair.

■ ■ ■ Medication

- Nonsteroidal anti-inflammatory medications, such as aspirin and ibuprofen (do not take within 7 days before surgery), or other minor pain relievers, such as acetaminophen, are often recommended. Take these as directed. Contact your physician immediately if any bleeding, stomach upset, or signs of an allergic reaction occur.
- Pain relievers are usually not prescribed for this condition. If your physician prescribes pain medications, use only as directed.
- Cortisone injections reduce inflammation. However, these are done only in extreme cases because there is a limit to the number of times cortisone may be given; it may

weakened muscle and tendon tissue. Anesthetics temporarily relieve pain.

■ ■ ■ Cold Therapy

Cold is used to relieve pain and reduce inflammation. Cold should be applied for 10 to 15 minutes every 2 to 3 hours for inflammation and pain and immediately after any activity that aggravates your symptoms. Use ice packs or an ice massage.

■ ■ ■ Notify Our Office If

- Symptoms get worse or do not improve in 2 to 4 weeks despite treatment
- You experience pain, numbness, or coldness in the hand
- Blue, gray, or dusky color appears in the fingernails
- Any of the following occur after surgery: increased pain, swelling, redness, drainage, or bleeding in the surgical area or signs of infection
- New, unexplained symptoms develop (drugs used in treatment may produce side effects)




Figure 1

From Hislop HJ, Montgomery J: Daniels and Worthingham's Muscle Testing—Techniques of Manual Examination, 6th ed. Philadelphia, WB Saunders, 1995, p. 152.

Notes:

(Up to 4400 characters only)

Notes and suggestions